

Germination and vigor of lettuce seeds (*Lactuca sativa* L.) pelleted with homeopathic preparations *Alumina* and *Calcarea carbonica* subjected to toxic levels of aluminum.

Filipe Pereira Giardini Bonfim¹, Rosana Gonçalves Rodrigues das Dores²,
Ernane Ronie Martins³, Vicente Wagner Dias Casali¹

(1) Federal University of Viçosa, MG, Brazil

(2) Federal University of Ouro Preto, MG, Brazil

(3) Federal University of Minas Gerais, Montes Claros, MG, Brazil

ABSTRACT

Background: aluminum toxicity is the most important factor limiting the growth of plants in acid soils, whereas current treatments are unfeasible. For this reason, alternatives are sought for, among which homeopathic treatment. **Aims:** this study aimed at evaluating the influence of homeopathic preparations *Alumina* 6cH, *Alumina* 12cH, *Calcarea carbonica* 6cH and *Calcarea carbonica* 12cH on the germination and vigor of lettuce seeds subjected to toxic levels of aluminum in paper-solution. At the same time, it was sought to develop a new procedure to apply homeopathic preparations in plants (pelleting). **Methods:** the statistical design was entirely randomized (CRD) with 6 treatments and 4 repetitions. Treatments included: 1) pelleted seeds/talc + *Alum* 6 cH; 2) pelleted seeds/talc + *Alum* 12cH; 3) pelleted seed/talc + *Calc* 6cH; 4) pelleted seeds/talc + *Calc* 12cH; 5) pelleted seeds/talc + distilled water; 6) non pelleted seeds (control). Variables evaluated were: germination percentage (GP), germination speed index (GSI) and radicle length (RL). **Results:** there was significant difference in GSI and RL – variables that reflect the vigor of seeds - between the samples treated with homeopathic preparations and the controls **Conclusions:** homeopathic preparations *Alumina* 6cH and 12cH and *Calcarea carbonica* 6cH and 12cH had significant effect on the vigor of lettuce seeds subjected to stress conditions.

Keywords: Aluminum toxicity; Homeopathic preparations; Lettuce; Germination; Vigor; Pelleted seeds

Introduction

Aluminum toxicity is the most important limiting factor for the growth of plants in acid soils [1]. It is held that the main effect of toxic levels of aluminum is a reduction in the rate of radicular growth of sensitive plants, since they affect the lengthening and division of cells [2]. This restriction decreases the plant's ability to take water and nutrients from the subsoil due to superficial rooting; in this way, it becomes less productive and more susceptible to drought.

To attempt to decrease this toxic effect in such soils correctives (calcareous compounds) and fertilizers are added in-depth. Nevertheless, the techniques currently available for this goal are unfeasible, partly because there is not yet a method to control the exchangeable aluminum at the sub-superficial level of soils, partly due to the cost of correctives and finally, and partly due to the large areas of soils presenting significant degrees of damaging acidity [3]; furthermore, these resources are not renewable, thus, they can be exhausted. For these reasons, new technologies are sought for.

Homeopathy is one among such technologies; it was approved in Brazil by the Ministry of Agriculture and Supply in 2008 [4]. The use of diluted and agitated substances in agriculture, especially in plants, is spreading fast. Notions and methods proper to homeopathy are used in several aspects of agriculture, including the quality of seedlings [5], germination [6], plagues control, [7] plants' diseases [8], increase of active principles [9], metal detoxification [10], and plant metabolism [11].

Homeopathy has potential to contribute to sustainable agriculture, this is the management of resources in order to satisfy the ever-changing human needs and at the same time keep or improve the quality of the environment and conserve the natural resources [12]. Through self-regulation, plants respond with high intensity to homeopathy when employed in stress conditions [13].

Despite positive results, both in the academic and practical levels, not too much is known regarding the physiological mechanisms of action of homeopathic preparations in plants, nor about the methods to employ them.

This study seeks to contribute to the development of a technology to facilitate the germination of plants in poorly favorable environments, as well as to the introduction of an innovative method to employ homeopathic preparations, such as seeds pelleting. In this context, it was sought to evaluate the influence of homeopathic preparations *Alumina* and *Calcarea carbonica* as seed pellets on the germination and vigor of seeds of lettuce subjected to toxic levels of aluminum in paper solution.

Materials and Methods

Choice of the homeopathic preparations

Homeopathic preparations *Alum* 6cH, *Alum* 12cH, *Calc* 6cH and *Calc* 12cH were selected according their indications in plants [14]. *Alumina* is indicated in slow-growing plants; growth interrupted due to excessive absorption of aluminum; culture in soils rich in aluminum leading to the accumulation of toxic levels; culture in acid soils. *Calcarea carbonica*: indicated in plants sensitive to adverse conditions; slow production of roots; slow growth with late sprouting, slow germination, smaller seeds and higher rate of sterility.

Preparation of the homeopathic dilutions

Agitated dilutions of *Alumina* and *Calcarea carbonica* were prepared from homeopathic matrixes in dilution 5cH (Schraiber) obtained at a homeopathic pharmacy in Viçosa, Minas Gerais, Brazil in October, 2009; further dilutions were prepared at the Homeopathic Laboratory of the Department of Phytotechnics at Federal University of Viçosa (DFT/UFV) according to the guidelines in the Brazilian Homeopathic Pharmacopoeia [15]. Succussion was carried out through a "mechanic-arm" device model DENISE 10-50 (UTIC). Homeopathic solutions *Alum* 6cH and 12cH, and *Calc* 6cH and 12cH were prepared immediately before the pelleting of seeds.

Seeds

For this study it was used naked lettuce seeds brand Isla ® type “Regina de Verão”, batch #22419 proceeding from Chile; germination percentage 91%; purity: 99.4%.

Pelleting of seeds

To pellet the seeds we used our own method of impregnation, observing the weight/volume ratio, as 1 gram of powder per ml of adhesive. Powder was Synth® inert pharmaceutical talc ($Mg_3Si_4O_{10}(OH)$); and adhesive corresponded to homeopathic preparations *Alum 6cH*, *Alum 12cH*, *Calc 6cH*, *Calc 12cH* and distilled water. The latter were made impregnate the talc in a Petri dish diving rise to a semi-pasty, integral and white powder, which was later used to coat the seeds.

Determination of the toxic dose of aluminum

A pretest was carried out to establish the concentration of Al^{3+} that best made patent its toxic effect on seeds of lettuce (*Lactuca sativa* L.) through the method of paper-solution [16]. In this sense, the best behavior was observed with a concentration twice the tolerable levels of lettuce cultures, which bears about 3.0 cmmol/dm^3 of Al^{3+} [17]. Consequently, it was used a solution corresponding to 540 mg/l^{-1} of $AlCl_3 \cdot 6H_2O$.

Experimental Design

The study was carried out at the Ultradilutions Laboratory of DFT/UFV, geographic coordinates $42^\circ 52'W$ and $42^\circ 50'W$ (longitude), and $20^\circ 44'S$ and $20^\circ 47'S$ (latitude). The statistical designed was entirely randomized (CRD), with 6 treatments and 4 repetitions; each experimental unit comprised 25 seeds. Treatments were: 1) pelleted seeds/talc + *Alum 6cH*; 2) pelleted seeds/talc + *Alum 12cH*; 3) pelleted seeds/talc + *Calc 6cH*; pelleted seeds/talc + *Calc 12cH*; pelleted seeds/talc + distilled water (control 1); non-pelleted seeds (control 2).

All seeds were placed in a germination chamber Tecnal® model TE-401in gerbox boxes with germitest paper previously moistened by 8 ml of aluminum solution (540 mg/l^{-1} of $AlCl_3 \cdot 6H_2O$), kept at $20^\circ C$, photoperiod 16 hours light and 8 hours dark, for 7 days. The tests of germination and vigor followed the guidelines and criteria established by the Ministry of Agriculture [18]. The experiment was carried out in a double-blind design, using symbols to name each treatment in order to avoid potential interferences.

Evaluated variables

Variables assessed included: germination percentage (GP), radical length (RL); germination speed index (GSI). Data were subjected to variance analyses and means were compared through Tukey's test with 5% probability with software SAEG. GP was established on the 7th day after seeding, counting the total number of seeds in each lot; GP data were transformed into $\sin^{-1} \sqrt{X/100}$. RL was measured at the end of the study with a digital paquimeter, and the results were expressed in millimeters (mm). GSI was carried out following Maguire [19], as the addition of the ratio between the number of germinated seeds each day and the day of evaluation: $IVG = G_1/N_1 + G_2/N_2 + G_3/N_3 + \dots + G_n/N_n$, where $G_1, G_2, G_3, \dots, G_n$ = number of germinated seeds in the day of observation, and $N_1, N_2, N_3, \dots, N_n$ = the number of days after seeding.

Results and Discussion

There were significant differences among treatments regarding the variables related to the vigor of seeds (GSI and RL), whereas there was no difference regarding GP, as shown in Table 1.

Table 1. Summary of variance analysis of germination percentage (GP), germination speed index (GSI) and radicle length (RL) in seeds of lettuce non-pelleted (control) and pelleted (with homeopathic preparations *Alumina* and *Calcarea carbonica*) subjected to toxic levels of aluminum.

Variation source	Degree of freedom	Mean squares		
		GP	GSI	RL
Treatments	5	65,86 ^{ns}	17,01 ^{**}	445,32 ^{**}
Residues	18	88	0,83	4,69
Total	23			
Variation coefficient (%)		10,99	8,29	11,65

(**) Significant at 1% probability respectively with f-test.

(ns) Non significant at 5% probability with f-test.

Treatment 5 (control 1) did not differ from non-pelleted Treatment 6 (control 2) in all 3 evaluated variables (Tables 2, 3) showing thus that the impregnating substance (pharmaceutical talc) had no interference effects between both controls and, thus, that the pelleting of seeds by itself, without homeopathic preparations, produces no benefit.

Figures 1 and 2 illustrate the germination and vigor of lettuce seeds on the fourth day and seven days respectively after the mounting of the experiment.




Figure 1 – Lettuce seeds, paper solution with toxic levels of aluminum (4th day): A) Treatment 3 (pelleted seeds/talc+*Calc* 6cH); B) Treatment 4 (pelleted seeds/talc+*Calc* 12cH); C) Treatment 6 – control 2 (non-pelleted seeds); D) Treatment 1 (pelleted seeds/talc+*Alum* 6cH); E) Treatment 2 (pelleted seeds/talc+*Alum* 12cH); F) Treatment 5- control 1 (pelleted seeds/talc+distilled water).

Regarding RL, Table 2 shows the significantly higher means of Treatments 1, 2, 3 and 4 by comparison with both control treatments, therefore, the effectiveness of low dilutions of *Alumina* and *Calcarea carbonica* on the physiological aspects of the seeds of lettuce. These results corroborate previously obtained ones [20], where it was observed an increase in the weight of the dry mass of lettuce treated with homeopathic preparation *Arnica montana* in dilutions 6cH and 12cH, which was not found with dilution 30cH.




Figure 2 – Small plants resulting from this study, final evaluation (7th day). A) Treatment 3 (pelleted seeds/talc+*Calc* 6cH); B) Treatment 4 (pelleted seeds/talc+*Calc* 12cH); C) Treatment 6 – control 2 (non-pelleted seeds); D) Treatment 1 (pelleted seeds/talc+*Alum* 6cH); E) Treatment 2 (pelleted seeds/talc+*Alum* 12cH); F) Treatment 5- control 1 (pelleted seeds/talc+distilled water).

Table 2 – Mean values of radicle length (RL) of lettuce seeds both non-pelleted and pelleted with homeopathic preparations of *Alumina* and *Calcarea carbonica* subjected to toxic levels of aluminum.

Treatments	RL (mm)
1. Pelleted seeds / Talc + <i>Alum</i> 6cH	25,43 a
2. Pelleted seeds / Talc + <i>Alum</i> 12cH	24,34 a
3. Pelleted seeds/ Talc + <i>Calc</i> 6cH	25,14 a
4. Pelleted seeds / Talc + <i>Calc</i> 12cH	24,61 a
5. Pelleted seeds / Talc + distilled water (control 1)	5,62 b
6. Non-pelleted seeds (control 2)	4,40 b

Means followed by at least one small-case letter in the line do not differ significantly between them in Tukey's test at 1% probability.

Regarding the germination speed index (GSI), Table 3 shows that the response of Treatments 1, 2, 3, 4 was similar to RL: all four differed from both controls; *Alum* 6cH and *Calc* 6cH obtained the highest means. This shows a positive effect of these homeopathic preparations on seeds exposed to unfavorable conditions. A similar behavior was seen in plants of *Ocimum basilicum* L. (basil) intoxicated with copper sulfate; following treatment with *Cuprum metallicum* 30cH drastically reduced the effect of the intoxication on the development of the plants.

Table 3 – Mean values of germination speed index (GSI) of lettuce seeds both non-pelleted and pelleted with homeopathic preparations of *Alumina* and *Calcarea carbonica* subjected to toxic levels of aluminum

Treatments	GSI
1. Pelleted seeds / Talc + <i>Alum</i> 6cH	12.38 a
2. Pelleted seeds / Talc + <i>Alum</i> 12cH	11.82 a
3. Pelleted seeds / Talc + <i>Calc</i> carb 6cH	13.05 a
4. Pelleted seeds / Talc + <i>Calc</i> carb 12cH	11.85 a
5. Pelleted seeds / Talc + distilled water (control 1)	8.17 b
6. Non-pelleted seeds (control 2)	8.60 b

Means followed by at least one small-case letter in the line do not differ significantly between them in Tukey's test at 1% probability.

Conclusions

This study showed that elleting, or the process of coating seeds, with homeopathic preparations represents an effective alternative means to employ such dilutions in plants.

Homeopathic preparations *Alumina* 6cH and 12cH, and *Calcarea carbonica* 6cH and 12cH had significant positive effects on the seeds of lettuce exposed to toxic levels of aluminum.

References

- [1] Lopes CJ, Capucho MT, Zanotti P. Germinação de sementes de espécies florestais de *Caesalpineia ferrea* Mart. Ex Tul. Var. *leiostachya* Benth., *Cassia grandis* L. E *Samanea saman* Merrill, após tratamento para superar a dormência. Revista Brasileira de Botânica. 1998; 20:80-86. [portuguese].
- [2] Kochian LV. Cellular mechanism of aluminium toxicity and resistance in plants. Annual Review of Plant Physiology. 1995; 46: 237-260.
- [3] Fortunato RP, Nicoloso, FT. Toxidez de alumínio em plântulas de grápia (*Apuleia leiocarpa* Vog. Macbride). Ciência Rura. 2004; 34(1):89-95. [portuguese].
- [4] Brasil. Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa nº 64, de 18 de dezembro de 2008. Aprova o Regulamento Técnico para os Sistemas Orgânicos de Produção Animal e Vegetal. Diário Oficial [da] República Federativa do Brasil, Brasília, DF, 19 de dezembro de 2008, Seção 1, p. 21.

- [5] Bonfim FPG, Martins ER, Dores RGR, Barbosa CKR, Casali VWD, Honório ICG. Use of homeopathic *Arnica montana* for the issuance of roots of *Rosmarinus officinalis* L. and *Lippia alba* (Mill) N.E.Br. Int J High Dilution Res [online]. 2008 [cited 2010 Aug 17]: 7(23); 113-117. Available from: <http://www.feg.unesp.br/~ojs/index.php/ijhdr/article/view/276/348> .
- [6] Hamman B, Konning G, Lok KL. Homeopathically prepared giberellic acid and barley seed germination. Homeopathy. 2003;92(3):140-144.
- [7] Almeida AA, Galvão JCC, Casali VWD, Lima ER, Miranda GV. Tratamentos homeopáticos e densidade populacional de *Spodoptera frugiperda* (J.E.Smith,1797) (*Lepidoptera: Noctuidae*) em plantas de milho no campo. Revista Brasileira de Milho e Sorgo. 2003; 2(2):32-39. [portuguese].
- [8] Khanna KK, Chandra S. Control of tomato fruit rot caused by *Fusarium roseum* with homeopathic drugs. Indian Phytopathology. 1976;29: 269-272.
- [9] Andrade FMC, Casali VWD, De Vita B. Efeito de homeopatas no crescimento e na produção de cumarina em chambá (*Justicia pectoralis* Jacq.). Revista Brasileira de Plantas Mediciniais.2001; 4, 1:19-27. [portuguese].
- [10] Almeida MAZ. Casali VWD, Cecon PR. Resposta do manjericão (*Ocimum basilicum* L.) à aplicação de preparações homeopáticas [dissertation]. Viçosa: Federal University of Viçosa; 2002.
- [11] Arruda VM. Aplicações de soluções homeopáticas em *Achillea millefolium* L. (Asteraceae): abordagem morfofisiológica [dissertation]. Viçosa: Federal University of Viçosa; 2005. [portuguese].
- [12] Lisboa SP, Cupertino MC, Arruda VM, Casali VWD. Nova visão dos organismos vivos e o equilíbrio pela homeopatia. Viçosa(Brazil):UFV; 2005. [portuguese].
- [13] Casali VWD. Utilização da homeopatia em vegetais. In: Proceedings of the Seminário Brasileiro sobre Homeopatia na Agropecuária Orgânica; 2004; Toledo, Brazil; 2004, 89-117. [portuguese].
- [14] Casali VWD, Andrade FMC, Duarte ESM. Ecologia de Altas Diluições. Viçosa(Brazil):UFV; 2009. [portuguese].
- [15] Farmacopéia Homeopática Brasileira. 1st ed. São Paulo (Brazil):Andrei; 1977. [portuguese].
- [16] Konzak CF, Polle E, Kittrick JA. Screening several crops for aluminum tolerance. In: Workshop on plant adaptation to mineral stress in problem soils; 1976; Beltsville. Ithaca: Cornell University Press; 1976. p.311-327.
- [17]. Ribeiro AC, Guimarães PTG, Venegas VHA. Recomendações para o uso de corretivos e fertilizantes em Minas Gerais: 5ª aproximação. Viçosa:CFSEMG; 1999. [portuguese].
- [18] Brasil. Ministério da Agricultura, do Abastecimento e da Reforma Agrária. Regras para análise de sementes. Brasília, DF(Brazil);1992. [portuguese].
- [19] Maguire J. Speed of germination-aid in selection and evaluation for seedling emergence and vigor. Crop Science. 1962; 2(1): 176-177.
- [20] Grisa S, Toledo MV, Oliveira LC, Holz L, Marine D. Crescimento e produtividade de alface sob diferentes potências do medicamento homeopático *Arnica montana*. Rev. Bras. de Agroecologia. 2007; 2: 1050-3.

Germinação e vigor de sementes de alface (*lactuca sativa* L.) Peletizadas com as preparações homeopáticas *alumina* e *calcarea carbonica* expostas a níveis tóxicos de alumínio.

RESUMO

Introdução: a toxicidade por alumínio é o principal fator que limita o crescimento das plantas em solos ácido; no entanto, os tratamentos disponíveis não são praticáveis. Por esse motivo, são procuradas alternativas, entre as quais, encontra-se a homeopatia. **Objetivos:** avaliar a influência das preparações homeopáticas *Alumina* 6cH e 12cH, e *Calcarea carbonica* 6cH e 12cH na germinação e vigor de sementes de alface expostas a níveis tóxicos de alumínio em papel-solução. Ao mesmo tempo, procurou-se desenvolver um novo método de aplicação de preparações homeopáticas nas plantas (peletização). **Métodos:** o desenho estatístico foi totalmente casualizado, com 6 tratamentos e 4 repetições. Os tratamentos incluíram: 1) sementes peletizadas/talco + *Alum* 6cH; 2) sementes peletizadas/talco + *Alum* cH12; 3) sementes peletizadas/talco + *Calc* 12cH; 4) sementes peletizadas/talco + *Calc* 12cH; 5) sementes peletizadas/talco + água destilada (controle 1) e 6) sementes não peletizadas. As variáveis avaliadas foram: porcentagem de germinação; índice de velocidade de germinação (IVG); e crescimento radicular (CR). Resultados: houve diferença significativa nas variáveis que refletem o vigor das sementes (IVG e CR) entre as amostradas tratadas com preparações homeopáticas e os controles. Conclusões: as preparações homeopáticas *Alumina* 6cH e 12cH, e *Calcarea carbonica* 6cH e 12cH apresentaram efeito significativamente positivo no vigor de sementes de alface expostas a condições de estresse.

Palavras-chave: Toxicidade por alumínio; Homeopatia; Alface; Germinação; Vigor; Sementes peletizadas

Germinación y vigor de semillas de lechuga (*lactuca sativa* L.) peletizadas con los preparados homeopáticos *Alumina* y *Calcarea carbonica* expuestas a niveles tóxicos de aluminio

RESUMEN

Introducción: la toxicidad por aluminio es el factor más importante que limita el crecimiento de las plantas en suelos ácidos, mientras que los tratamientos disponibles son impracticables. Por eso, se buscan alternativas, entre ellas, la homeopatía. **Objetivos:** evaluar la influencia de los preparados homeopáticos *Alumina* 6cH y 12cH, y *Calcarea carbonica* 6cH y 12cH en la germinación y vigor de semillas de lechuga expuestas a niveles tóxicos de aluminio en papel-solución. Al mismo tiempo, se buscó desarrollar un nuevo método para aplicar preparados homeopáticos en plantas (peletización). **Métodos:** o diseño estadístico fue completamente casualizado, con 6 tratamientos y 4 repeticiones. Los tratamientos fueron: 1) semillas peletizadas/talco + *Alum* 6cH; 2) semillas peletizadas/talco + *Alum* cH12; 3) semillas peletizadas/talco + *Calc* 12cH; 4) semillas peletizadas/talco + *Calc* 12cH; 5) semillas peletizadas/talco + agua destilada (controle 1) e 6) semillas no peletizadas. Las variables evaluadas fueron: porcentaje de germinación (PG); índice de velocidad de germinación (IVG); y longitud de las radículas (LR). **Resultados:** hubo diferencia significativa en el IVG y LR – que reflejan el vigor de las semillas – entre las semillas tratadas con preparados homeopáticos y los controles. **Conclusión:** los preparados homeopáticos *Alumina* 6cH y 12cH, y *Calcarea carbonica* 6cH y 12cH manifestaron efectos estadísticamente significativos en el vigor de semillas de lechuga expuestas a condiciones de stress.

Palabras-llave: Toxicidad por aluminio; Preparados homeopáticos; Lechuga; Germinación; Vigor; Semillas peletizadas.



Licensed to [GIRI](#)

Support: CNPq

Conflict of interest: authors declare there is no conflict of interest

Received: 16 August 2010; Revised: 05 December 2010; Published: 20 December 2010.

Correspondence author: Filipe Pereira Giardini Bonfim, filipegiardini@yahoo.com.br, Departamento de Fitotecnia da Universidade Federal de Viçosa. Av. PH. Rolfs, s/n. Campus Universitário. Viçosa. MG. Brasil. CEP: 36571000.

How to cite this article: Bonfim FPG, das Dores RGR, Martins ER, Casali VWD. Germination and vigor of lettuce seeds (*Lactuca sativa* L.) pelleted with homeopathic preparations *Alumina* and *Calcarea carbonica* subjected to toxic levels of aluminum. Int J High Dilution Res [online]. 2010 [cited YYYY Month dd]; 9(33): 138-146. Available from: <http://www.feg.unesp.br/~ojs/index.php/ijhdr/article/view/412/450>.