Abstract

Homeopathy and Levels of Ca⁺⁺ Salivary in the Prevention of Dental Caries

Leila C. Mourao*, Rosy Nardy, Dionisia Cataldo, Vicente Canuto, Adriana Marques Nunes, Leonardo Barroso, Aisha Vassem, Irene Cunha Ramos, Kadija Bonaud, Erick Santos, Thamires Lorena

UniFoa – Centro Universitário Volta Redonda

*leilamourao@hotmail.com - <https://orcid.org/0000-0002-8682-7580>

Comparative studies between systemic inducing bone neoformation medications showed favorable results for homeopathic medicines. (Almeida et al., 2000; Werkman et al., 2006, FARINA, 2010). The Calcareas, are described in the literature for their action in the mineralization processes and increase of the levels of Ca ++ tissue ((LATHOUD, 1998; ARGENTA, 2005; VIJNOVSKY, 1999; VOISIN, 1971). NARDY-MELO, 2013, found that Calcareas carbonica 6CH, Calcareas phosphorica 6CH and Calcareas fluorica 6CH, when given as a compound, are statistically significant for increasing tissue Ca ++ levels. **Objective** This randomized controlled multicentric study, approved by the UniFOA's Human beings Research Ethics Committee CAAE:39002914.0.0000.5237, has aspiration to quantify the level of Ca ++salivary in front of the administration of homeopathic medicines Calcareas Carbonica, Calcareas Phosphorica, Calcareas Fluorica, as supporting in the prevention of dental caries, Demineralization and Dental I, from the stimulus to increase of Ca ++ salivary. **Method** (n=48) Children in the age range of 04 to 16 years, with high rate of dental caries, submitted to dental treatment, without the preventive application of fluoride topical and specific food guidelines were divided randomly, in 2 groups according to the medication experienced: GI – *Calcareas Carbonica* 6CH; *Calcareas phosphorica* 6 CH; *Calcareas fluorica* 6CH, in the dose of 3 daily drops for 15 days a month for three months. GII – Upward scales of *Calcareas Carbonica* 6CH; *Calcareas phosphorica* 6CH; *Calcareas fluorica* 6CH, in the dose of 3 daily drops, in the first month. *Calcareas Carbonica* 9CH; *Calcareas phosphorica* 9CH; *Calcareas fluorica* 9CH, in the dose of 3 daily drops, in the second month. *Calcareas Carbonica* 12CH; *Calcareas phosphorica* 12CH; *Calcareas fluorica* 12CH, in the dose of 3 daily drops, in the third month. The salivary Ca + + levels was measured before the administration of the first dose and monthly, for 03 consecutive months. The calcium (mg/dl) salivary indices were determined by optical emission spectrometry (ICP-OES). **Results** The results were submitted to statistical analysis with Student T-Test (significance with P < 0.05) and analysis of variance Shapiro-Wilk normality test. **Conclusion:** The averages of calcium increase under the action of homeopathic medicinal products: *Calcareas Carbonica*, *Calcareas Phosphorica* and *Calcareas Fluorica*, in this study, do not differ in G I and G II and in both groups the increase in salivary calcium level was statistically significant to 0.5%.

Keywords: homeopathy, salivary calcium, prevention

© International Journal of High Dilution Research.
Not for commercial purposes.