Prospective study of ADHD cases with biological confirmation

Fabio de Almeida Bolognani

Federação Brasileira de Homeopatia, Rio de janeiro, Brazil

Introduction: Based on the large amount of patients that seek alternative treatment for clinical complaints compatible with ADHD diagnosis and above all by the high percentage of non adaptation or risk that officially proposed drugs may bring, we developed a laboratorial clinical investigative profile that favors a safe homeopathic treatment.

Material: 52 cases with urinary zinc analysis, with ADHD diagnosis.

Method: using the diagnosis confirmation questionnaire (SNAP-IV by UFRJ) and Zinc analysis by HPLC method (high performance liquid chromatography) which normal values range from 246 and 846 nmol/liter.

Results: in a 52 patients (table 1) sampling with a suggestive ADHD diagnosis, we obtained a quantitative alteration in 31 cases showing high urinary Zinc levels, which allowed therapeutic approach that lead to better clinical result followed by subsequent reduction of urinary Zinc levels, in posterior analysis in eight (8) of these patients.

Table 1: 52 patients (table 1) sampling with a suggestive ADHD diagnosis

	Standard	1st exam	2nd exam	3rd exam	4th exam
	266 / 846mcl / l				
1-Ayslan		1625	1211	2007	719
2-Annie		3203			
3-Bernardo		1113			
4-Bruna		669			
5-Bernardo S		1061			
6-Erik		527.8			
7-Estefania		3463			
8-Gabriel		615			
9-Gabriela		319,1			
10-Italo		356.9			
11-Ian Jose		1442	181.1	309,5	
12-Igor		340.8			
13-Iann		588			
14-Isabel		1836	646		
15-Joao Lor		982			
16-Kaleb		3871	2956	5590	378
17-Leonora		2319			
18-Lara		4704	2055	3424	1338
19-Michel		857.9			
20-Murilo		7002	435		
21-Matheus		1400	3360	704.5	594
22-Maria Clara		1265			
23-Mateus		453			
24-Paulo Henri		974			
25-Pedro		334			
27-Nycole		1004	905	484	
28-Pedro Ab		433	166		

29-Rodrigo Ch		1115			
30-Rodrigo Ho		820			
31-Rodrigo Lim		965	800		
32-Thiago Kres		212			
33-Thales		1113	870	745	
34-Thiago		413			
35-Warnei		328.8			
36-Waldemar		1059			
37-Thiago Lour		560.7	927	1585	
38-Joao Vitor		884,1			
39-Pedro Henrique		1462,2			
40-Ana Carolina Jardim		970	755		
41-Luan		456,3			
42-Laura Merlim		1217,3			
43-Joao Pedro Papa		282			
44-Francisco V B Soares		436,3	370,2		
45-Thales Martoreli		1424	256		
46-Daniel Alves		828			
47-Luiz Eduardo The		943			
48-Daniel Silva		614,2			
49-Wagner Jardim		1091			
50-Gustavo Maciel		1133			
51-Matheus Holanda		1500	590	1100	
52-Gabriel Oliveira		1551			

Clinical results obtained in groups showing alteration in urinary zinc associated with subsequent evaluations, where introduction of *Zincum metallicum* medication, at 12 CH, led to reduction in zinc values in urine and consequently clinical amelioration.

Other patients undergoing exams, which weren't included in subsequent evaluations, showed improvement, but since exams were executed in different laboratories other than the laboratory from the first analysis, weren't considered.

Patients numbers: 1, 16 and 18 (table 2) demonstrated a raise in urinary zinc levels from second to third analysis. These patients stopped using their homeopathic medications for at least 30 days (during school vacation), which lead to this increase but when reintroducing the medication, these rates lowered to normal levels again.

Table 2: Patients with a raise in urinary zinc levels from second to third analysis.

18-Lara		4704	2055	3424	1338
16-Kaleb		3871	2956	5590	378
1-Ayslan		1625	1211	2007	719

Comments:

Treatment of this clinical condition places the homeopathic doctor in different situations according to legislations and rules for medical procedures.

Some countries consider that from Attention Deficit Hyperactivity Disorder diagnosis, the use of specific medications is mandatory, and its use will be restricted only in cases where its use leads to harm.

Homeopathic doctors also encounter difficulties with these patients, since medications even while being appropriate; do not always offer safety of lasting clinical results.

Alterations in human body led by Zinc element, when unstable, due to environmental phenomena, could justify the concept of “epidemic gene” since symptoms and origins demonstrate susceptible population of considerable proportion.

Attention to hygiene and food habits are the main issues approached, but habits and food availability in each population can determine incidence and re-incidence of this clinical phenomenon. Why?

People without diagnose of TDAH

1-	Suelene Dias Silva	153
2-	Leo Pereira Silva	485
3-	Erik Pereira	760,1
4-	Angelo Pereira	302
5-	Angela Bazzanela	736
6-	Sonia Palombini	264
7-	Paulo C Gomes	502,7
8-	Larissa Primo	555
9-	Lara Salles	332
10-	Robert Vieira	670,6

Zincum Mettalicum:

- in metabolism of over 100 enzymes, and interferes in hormone functioning, protein synthesis, reproduction and immunological system.
- Recent article, published in Nature Reviews, Neuroscience, November 2009/ volume 10, by Sensi (University G. d’Annunzio-Italy and University of California-USA), Paoletti (Centre National de Recherche-Paris-France), Bush (University of Melbourne-Australia) and Sekler (Ben Gurion University-Israel), reveals that in the brain, zinc in its ionic form (Zn^{2+}) is highly enriched at many but not all glutamatergic nerve terminals, is released upon neuronal activity and is crucial to the control of physiological and pathophysiological brain functions.

This Zinc, is estimated to rise to micromolecular levels in the proximity of axon terminals following release from synaptic vesicles that contain millimolar (Zn^{2+}).

Synaptically released Zn^{2+} interacts with various neuronal ion channels, receptors and transporters and therefore modulates synaptic transmissions and plasticity.

Zn^{2+} affects the activity of NMDA (N-methyl-D-aspartate) and GABA (γ -amino butyric acid type A) receptors, thereby influencing excitatory synaptic transmission in the brain.

Besides its physiological functions, Zn^{2+} is also a potent toxin that is involved in neuronal and glial death and is associated with a variety of excitotoxic conditions, such as epilepsy, ischaemic and brain trauma. Finally, a major role for Zn^{2+} has been suggested in the neurodegeneration associated with Alzheimer’s disease, as it is a key component of amyloid plaques and the cerebral amyloid angiopathy observed in Alzheimer’s disease.

The best studied Zn transporter in the brain, is mainly localized at Zn^{2+} containing synaptic glutamatergic vesicles in the hippocampus, cortex and olfactory bulb, and is also present in certain cerebella GABAergic and

dopaminergic neurons and in specific locations of the spinal cord. Others studies reveals the fact that severe Zn²⁺ deficiency, caused by diet or chelation, enhances susceptibility to seizures.

The functional relevance of such sophisticated Zn²⁺ homeostatic machinery that controls the balance of Zn²⁺ concentrations between synaptic and somatic compartments of neurons is underlined by the emerging role of Zn²⁺ in the modulation of brain functioning.

Many glutamatergic vesicles in certain brain regions (neocortex, amygdale and hippocampus) contain high concentrations of Zn²⁺, suggesting that it has a role in cognition and synaptic plasticity. There are strong evidences that Zn²⁺ truly acts as a co-transmitter at many Central Nervous System synapses and participates in synaptic plasticity, and some kinds of experiences can show tremors, delayed righting reflex, abnormal gait and exaggerated startle responses in mice.

Further evidence for a modulatory role of Zn²⁺ in long term potentiation comes from recordings of hippocampus CA3 mossy fiber synapses, the most Zn²⁺ enriched synapses in the brain. At these synapses, extracellular Zn²⁺ may be present in a tonic fashion at low (probably nanomolar) levels, whereas levels can increase transiently during action potential driven exocytosis..

- It is widely present in pancreas, liver, skin and annexes.
- Some substances, such as PHYTATES, found in high fiber vegetables, as well PHYTALATES (polyvinyl chloride packages), alcohol, tannins, antibiotics (ampicillin), corticosteroids and contraceptives, inhibit dietary zinc absorption, or accelerates its elimination in conditions such as cirrhosis, renal illnesses, etc.

At this phase, we establish a relationship with Bisphenol-A, known as one of the most common environmental endocrine disrupter, that has been extensively evaluated for toxicity in a variety of tests in rodents, including developmental and reproductive toxicity, and carcinogenicity. However, in the report of Prenatal and neonatal exposure to Bisphenol-A, (Neuroscience 2003 / 117, by Suzuki, Nakazawa, Funae, Fukushima, Narita, from Departement of Toxicology, School of Pharmacy and Pharmaceutical Sciences-Tokio-Japan), show that prenatal and neonatal exposure to BPA in mice leads to enhancement of dopamine D1 receptor-dependent rewarding effect induced by a psycho stimulant methamphetamine. Furthermore, this treatment with BPA markedly enhanced hyper locomotion and its sensibilization induced by methamphetamine, which reflects extensive abuse associated with sociological and psychiatric problems.

In other report, Prenatal phthalate exposure and performance on the Neonatal Behavioral assessment scale, Neurotoxicology/2009/30, by Engel, Zhu, Berkowitz, Calafat, Silva, Wolff, from Mount Sinai School of Medicine-New York-USA), the relationship between prenatal urinary concentrations of phthalate metabolites and neonatal behavior in 295 children, that received by trained examiners the Brazelton Behavioral Assessment Scale. Seven phthalate esters metabolites were evaluated.

The result, among girls, there was a significant linear decline in adjusted mean Orientation score with increasing urinary concentrations of high molecular weight phthalate metabolites. Likewise, there was a strong linear decline in their adjusted mean Quality of Alertness score. In addition, boys and girls demonstrated opposite patterns association between low and high molecular weight phthalate concentrations and motor performance, with increasing concentration of low molecular weight phythalate metabolites among boys.

Janet Raloff, report in Science News, November/2009, the study of Bruce Lanphear, of Simon Fraser University in Burnaby-British Columbia. BPA in the womb shows link to kid's behavior, linked prenatal exposure to BPA with subtle, gender-specific alterations in behavior among two years olds. Girls whose mothers had encountered the most BPA early in pregnancy tended to become somewhat more aggressive

than normal, boys became more anxious and withdrawn. BPA, a common ingredient in hard polycarbonate plastics and resins used in food-can linings. Braun, in his paper in *Environmental Health Perspectives*, report that the higher a mom's BPA levels were during her first 16 weeks of pregnancy, the more likely her child was to later show behavior somewhat atypical of its gender. Braun report that the mannish girls and feminized boys, are more due a production or action of their gender-establishing sex hormones might have been blocked.

Girls whose mothers had the highest BPA exposures in early pregnancy tended to score high on the externalizing component of a test known as BASC-2, for Behavioral Assessment System for Children-2. It's geared to young children and evaluates their tendency towards aggression and hyperactivity.

Boys with the highest womb exposures to BPA tended to exhibit an increase internalizing BASC-2 score. They were more likely to be withdrawn or show depressive symptoms, or maybe complain about aches and pains, which is often a manifestation of anxiety.

- Neurological syndromes:
 - Physical asthenia, physical and psychic asthenia: depletion, dejection, and depression.
 - Intellectual, physical and sexual asthenia.
 - Slow comprehension, obnubilation,
 - Slow, but impatient and irritated
 - Intolerant to noises and conversations
 - Sensorial hyperesthesia
 - Insomnia with constant movement of legs and feet, fright, screams during sleep, wakes up frightened
 - Constant agitation of inferior members, especially when seated
 - When lays down to sleep, legs won't stop and prevents from sleeping
 - Enhances tendinous reflexes
 - Spasms, tremor, contractures, myoclonia
 - Convulsion, chorea, epilepsy
 - Lack of neuro muscular coordination
 - Urinary retention
 - Incontinence
 - All nervous pathologies, benign or chronic, following brutal retrogress or repression of infectious or eruptive disease, or associated with the use of corticosteroids, antibiotics, antidepressants, neuroleptic drugs, tranquilizers.

It would be possible that repulsion towards food from vegetal group, especially with more fibers that are common to children would be a form of "self" prophylaxis?

Food and beverage stored in polyvinyl packages, which is normally not perceived by basic human instincts, do not find the same resistance by human body?

Abuse of antibiotic and corticosteroid treatment usually seen today may take part in this alteration? Would alterations in digestive tract be important?

Meanwhile, results from this study suggest that association of *Zincum mettalicum*, might regulate what seems to be a greater loss of this element in urine, since blood analysis did not alter, which apparently corroborated the hypothesis of absorption of zinc being correct, reducing urinary loss.

Bibliography:

- ADAMS R., VICTOR M., ROPPER A., - Principles of Neurology, New York, International Edition, 1997.
- ALLEN, T.F. – The Encyclopedia of Pure Materia Medica, New Delhi. B. Jain Publishers, 990, vol II.
- ARIOVALDO, R.F. – Novo Repertório de Sintomas Homeopáticos, São Paulo. Robe Editorial, 1996.
- BLACKWOOD, A. L. – Materia Medica Terapeutica y Farmacologia Homeopatica. B. Jain Publishers.
- CONAN MERIADEC, L' Homeopathie, 1990, Paris, Editions Boiron
- CARILLO, R., Fundamentos da Homeopatia Constitucional, 1007, São Paulo, Santos editora
- CAIRO, N. – Guia de Medicina Homeopática, São Paulo. Livraria Teixeira, 1976.
- CLARCKE, J.H. – Dictionary of Practicas Materia Medica, New Delhi. Jain Publishing Co, 1983, vol II.
- COSTA, A.F. – Farmacognosia, Lisboa. Fundação Calouste Gulbenkian, 1987.
- DELLIERE, M., PASQUIER, A., Homeopathie et Sport, 2006, Marabout
- DEMARQUE, D., JOUANNY, J., POITEVIN, B., SAINT-JEAN, Y., Homeopathie connaitre la Matiere Medicale, Paris, 1989, CEDH
- FLEHMIG, I. – Desenvolvimento normal e seus desvios no lactente, Rio de Janeiro. Atheneu, 1982.
- GILMAN, A.G.; GOODMAN, L.S.; GILMAN, A. – As Bases Farmacológicas da Terapêutica, Rio de Janeiro. 1985.
- GUERMONPREZ, M. Matiere Medicale Homeopathique, Paris, 1985, Doin
- HAGER – Tratado de Farmácia Práctica, Barcelona. Editorial Labor, 1942, tomo II.
- HAHNEMANN, C.S.F. – Matéria Médica Pura, São Paulo. Editorial Homeopática Brasileira, 1998.
- HAHNEMANN, C.S.F. – Organon da Arte de Curar, São Paulo. GEHSP “Benoit Mure”, 1995.
- HAMILTON, E.M.D. – The Flora Homoeopathica, New Delhi. B. Jain Publishers, 1995.
- HERING, C.M.D. – The guiding Symptons of our Materia Medica, New Delhi. B. Jain Publishers, 1993, vol III.
- HODIAMONT, g., Homeopathie et Physiologie, 1983, Paris, J.B. Bailliers
- HUGLES, R. – A Manual of Pharmacodynamics, New Delhi. B. Jain Publishers, 1994.
- LUNDY-EKMAN, L., Neurociencia, 2002, EUA, Elsevier
- KENT, J.T. – Materia Medica Homeopatica, Buenos Aires. Editorial Aebatros, 1980, tomo II.
- KOLLITSCH, P. – Matière Médicale Therapeutique, Paris. Livrairie Maloene, 1955.
- NELSON – Textbook of Pediatrics, Philadelphia, London, Toronto. Vaughan and Mckay, 1975.

RHODE, MATTOS, Principios e Pratica em TDAH, Atmed Editora, 2003, Porto Alegre, Brasil

SCHIMDT, Neurofisiologia, 1979, São Paulo, Editora Pedagogica Universitaria

SENSI, PAOLETTI, BUSH, SEKLER, Zinc in the physiology and pathology of the CNS, Nature Reviews, November 2009, volume 10, Macmillan Publishers Limited

TETAU, M., Bergeret, C., L' Organotherapie diluee et dynamisee, Paris, 1984, Maloin

TYLER, M.L. – Retratos de Medicamentos Homeopáticos, São Paulo. Livraria Editora Santos, 1992.

VIJNOVSKY, B. – Tratado de Materia Medica Homeopatica, Buenos Aires. 1992, tomo II.

VOISIN, H. – Manual de Matéria Médica para o Clinical Homeopata, São Paulo. Organização Andrei Editora, 1984.

ZISSU, R.- Matiere Medicale Homeopathique Constitutionelle, Paris, 1977, Librairie Le François,

Licensed to [GIRI](#)

Support: authors declare that this study received no funding

Conflict of interest: authors declare there is no conflict of interest

Correspondence author: Fabio de Almeida Bolognani, fabibolognani@gmail.com , www.homeopatiabrasil.org.br

How to cite this article: Bolognani FA. Prospective study of ADHD cases with biological confirmation. Int J High Dilution Res [online]. 2011 [cited YYYY Month dd]; 10(35): 84-90. Proceedings of the XXIV GIRI Symposium; 2010 Nov 05; Monte Carlo (Monaco). GIRI; 2010. [cited YYYY Month dd]; Available from:

<http://www.feg.unesp.br/~ojs/index.php/ijhdr/article/view/443/475>