Conference presentation

Biological activity and phytochemical analysis of homeopathic preparations of *Viscum album* L.

Carla Holandino^{1*}, João V. da C. Batista¹, Michelle N. de O. Melo¹, Fortune Homsani¹, Adriana P. Oliveira¹, Leandro M. Rocha², Venício F. da Veiga³, Daniela S. Alviano³, Celuta S. Alviano³, Mirio Grazi⁴, Hartmut Ramm⁴, Matthias Kunz⁴, Konrad Urech⁴, Ulrike Weissenstein⁴, Stephan Baumgartner⁴

¹Multidisciplinary Laboratory of Pharmaceutical Sciences, School of Pharmacy, Federal University of Rio de Janeiro, Brazil; ² Fluminense Federal University, Niterói, Rio de Janeiro, Brazil; ³Institute of Microbiology, Federal University of Rio de Janeiro, Brazil; ⁴Society for cancer Research, Hiscia Institute, Arlesheim, Switzerland

*Corresponding author email: cholandino@gmail.com

Abstract

Several *in vitro* and *in vivo* studies investigated the biological properties of mistletoe preparations [1-4]. In Europe, where phytochemical constituents of aqueous *Viscum album* extracts were previously described, the most widespread and common form of mistletoe is the white-berried species (*Viscum album*) [5]. The aim of the present study was to analyze the biological activity and phytochemical features of different *Viscum album* ssp mother tinctures collected in 2 different harvesting periods.

Leaves, stems and berries of *Viscum album* samples were subjected to ethanolic extraction (45% v/v) following the homeopathic pharmacopeia [6,7]. The mother tinctures (MT) prepared were: *V. album* (from 3 host trees *Malus domestica*, *Quercus* sp, *Ulmus* sp), *V. album* ssp. *austriacum* (host tree *Pinus sylvestris*) and *V. album* ssp. *abietis* (host tree *Abies pectinata*). All samples were collected at Höfli farm (Switzerland) in July and August and subjected to chemical analysis by means of high-performance liquid chromatography-mass spectrometry (HPLC-MS/MS) and Thin-Layer Chromatography (TLC). Tumor (Yoshida and Molt4) and non-tumor cells (Ma104) were incubated with MT (1; 0.1; 0.01%) for 4, 24 and 48 hours at 37°C. Proliferation was indirectly measured using WST-1 (tumor cells) and MTT (non-tumor cells). All cells (1.5×10^4 /100 μ L) were plated in triplicate in 96-well plates. The cell proliferation inhibition rate was calculated as percent inhibition relative to control cells treated with the MT solvent (45% ethanol v/v). Apoptosis/necrosis was measured using 2×10^5 cells incubated with Annexin V-FITC and 7-AAD at room temperature in the dark, and analyzed in flow cytometer [2]. Antimicrobial activity was assessed for *Candida albicans*, *Cryptococcus neoformans*, *Escherichia coli* and *Staphylococcus aureus* following the Clinical Laboratory Standards Institute manuals [8-9] and compared with 45% ethanol v/v (MT solvent).

The chromatographic assays of MT samples showed phenolic acids and lignans as main chemical classes identified. WST-1 and MTT experiments revealed different antitumor potential of MT. In general, Yoshida cells were more sensitive than Molt4. The maximal cytotoxic effect with 100% of mortality was attained by incubation of Yoshida and Molt-4 cells with 1% MT of *V. album* ssp.

abietis (harvested in July) for 4 hours. Preliminary results with non-tumor cells indicate lower cytotoxicity compared to tumor cells. Annexin V/7ADD showed predominance of late apoptotic/necrotic events after 4 and 24 hours of incubation with all MT. The antimicrobial assay showed that MT of *Viscum album* ssp. *austriacum* (host tree *Pinus sylvestris*) exhibited minimum inhibitory concentration 7.25 mg/mL for the 2 tested bacterial strains.

The results of the present study suggest that *Viscum album* samples prepared according to the homeopathic pharmacopeia exhibit promising biological activities in a dose-dependent manner. The effects of potentiation will be assessed and compared to non-potentiated samples in future experiments. Therefore, the present study contributes with data on the main chemical components in homeopathic preparations of *Viscum album*, drawing attention to the relevance of extract solvent, seasonal aspects, host tree and harvested parts of plant harvested to the quality of the mother tinctures

Keywords: Mistletoe, *Viscum album* L. cancer, homeopathy, phytochemistry; cytotoxicity

References

- [1] Tröger W, Galun D, Reif M, Schumann A, Stankovic N, Milicévic M. *Viscum album* [L.] extract therapy in patients with locally advanced or metastatic pancreatic cancer: A randomised clinical trial on overall survival. *Eur J Cancer*. 2013;49(18):3788-3797.
- [2] Weissenstein U, Kunz M, Urech K, Regueiro U, Baumgartner S. Interaction of a standardized mistletoe (*Viscum album*) preparation with antitumor effects of trastuzumab in vitro. *BMC Complement and Altern Med*. 16:271, 2016, doi 10.1186/s12906-016-1246-2.
- [3] Mulsow K, Enzlein T, Delebinski C, Jaeger S, Seifert G, Meizig MF. Impact of mistletoe triterpene acids on the uptake of mistletoe lectin by cultured tumor cells. *Plos One* April 18, 2016, doi 10.1371/journal.pone.0153825.
- [4] Kang SN. Ethanol extracts from mistletoe (*Viscum album* L.) act as natural antioxidants and antimicrobial agents in uncooked pork patties during refrigerated storage. *Asian Australas. J. Anim. Sci*. 29: 109-118, 2016.
- [5] Urech K, Baumgartner S. Chemical constituents of *Viscum album* L.: implications for the pharmaceutical preparation of mistletoe. *Trans Res Biomed*. 2015; 4:11-23, doi 10.1159/000375422.
- [6] ANSM. Agence Nationale de Sécurité du Médicament et des Produits de Santé. Pharmacopée Française, Preparations-homeopathiques. The General Chapters and General Monographs of the European Pharmacopoeia and Preamble of the French Pharmacopoeia apply. Mistletoe from the apple tree for homeopathic preparations. 11e édition, Paris, 2011, pages 1-4.
- [7] ANVISA. Agência Nacional de Vigilância Sanitária. Farmacopeia Homeopática Brasileira, 3ª ed., Ed. ANVISA, Brasília; 2011; pages 31-33.
- [8] Clinical Laboratory Standards Institute (CLSI). Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically: Approved Standard, CLSI document M07-A10. 10. Ed. Wayne, PA: Clinical and Laboratory Standards Institute, 2015/<http://www.facm.ucl.ac.be/intranet/CLSI/CLSI-2015-M07-A10-original.pdf>, date of access: October 1st, 2017.

[9] Clinical and Laboratory Standards Institute (CLSI). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts: Approved Standard, CLSI document M27-A3. 3. Ed. Wayne, PA: Clinical and Laboratory Standards Institute, 2008/
https://clsi.org/media/1461/m27a3_sample.pdf, date of access: October 1st, 2017.

© International Journal of High Dilution Research/ Not for commercial purposes.